



Unlocking Epigenetic
Therapeutics to
Revolutionize Medicine

January 2025



Disclaimer

This presentation has been prepared by DURECT Corporation ("DURECT," the "Company," "we," "our" or "us") for informational purposes only. This document does not constitute an offer or invitation for the sale or purchase of securities. Neither the Company nor any of its affiliates makes any representation or warranty, express or implied, as to the accuracy or completeness of the information contained herein and shall have no liability for such information. To the fullest extent permitted by law, in no circumstances will the Company or its stockholders, affiliates, representatives, partners, directors, officers, employees, advisers or agents be responsible or liable for any direct, indirect or consequential loss or loss of profit arising from the use of this presentation, its contents, its omissions, reliance on the information contained within it, or on opinions communicated in relation thereto or otherwise arising in connection therewith. Interested parties should conduct their own investigation and analysis. The Company undertakes no obligation to provide the recipient with any additional information or to update or correct the information. The investment opportunity described herein is speculative and entails a high degree of risk.

This presentation and various remarks we make during this presentation contain forward-looking statements of DURECT within the meaning of applicable securities laws and regulations, which are subject to the safe harbor provisions of the Private Securities Litigation Reform Act of 1995, including statements relating to the potential for larsucosterol to demonstrate a reduction in mortality or liver transplant in patients with alcohol-associated hepatitis ("AH") and to save lives, DURECT's plans, trial design and preparation to initiate a 200-patient Phase 3 trial in the U.S. for larsucosterol with a 90-day survival endpoint with data expected within two years of trial initiation, the potential for a single Phase 3 trial of larsucosterol to show a statistically significant improvement in the treatment of AH over standard of care, the potential Federal Drug Agency (the "FDA") or other regulatory approval of larsucosterol for the treatment of AH after a Phase 3 trial, the potential to achieve over \$1 billion in peak sales in the U.S., anticipated physician and hospital awareness of AH and associated product benefits and other potential uses of larsucosterol, and anticipated product markets and incident projections. Actual results may differ materially from those contained in the forward-looking statements contained in this presentation, and reported results should not be considered as an indication of future performance. These forward-looking statements involve risks and uncertainties that could cause actual results to differ from those in such forward-looking statements. Potential risks and uncertainties include, but are not limited to, the risk that future clinical trials of larsucosterol are delayed or do not confirm the results from subset analyses of the AHFIRM trial, including geographic or other segmentation, or of earlier clinical or pre-clinical trials, or do not demonstrate the safety or efficacy of larsucosterol in a statistically significant manner, the risk that the FDA or other government agencies may require additional clinical trials for larsucosterol before approving larsucosterol for the treatment of AH, and that larsucosterol may never be approved; and risks related to the sufficiency of our cash resources, our anticipated capital requirements, our need or desire for additional financing including the need to obtain sufficient funding to initiate a Phase 3 trial for larsucosterol, our ability to regain compliance with the minimum bid price for continued listing on Nasdaq, our ability to obtain capital to fund our operations and expenses and our ability to continue to operate as a going concern. Further information regarding these and other risks is included in DURECT's most recent U.S. Securities and Exchange Commission ("SEC") filings, including its Annual and Quarterly Reports on Forms 10-K and 10-Q, respectively, filed with the SEC under the heading "Risk Factors." DURECT is under no duty to update any of these forward-looking statements after the date of this presentation to conform these statements to actual results or revised expectations, except as required by law. Existing and prospective investors are cautioned not to place undue reliance on these forward-looking statements, which speak only as of the date hereof and nothing (including any sale of securities by the Company after the date of this presentation) shall imply that the information contained herein or the affairs of the Company have not changed since the date hereof. Subsequent events and developments may cause DURECT's expectations and beliefs to change.

DURECT management based all estimates and projections upon their best judgment as of the date of this presentation and upon assumptions and circumstances that have not yet taken place, may not have an empirical basis, are subject to variation and are inherently unpredictable. There can be no assurance that any estimates or assumptions will prove accurate or that any of the projections will be realized. Actual results will vary from the projections, and such variations may be material.

This presentation is for informational purposes only and does not constitute an offer to buy or sell or a solicitation of an offer to buy or sell any securities of the Company. Any offer of securities will only be made pursuant to a registration statement (including a base prospectus) and prospectus supplement filed with the SEC, copies of which may be obtained for free on our website at www.durect.com under the "Investors" tab or by visiting EDGAR on the SEC website at www.sec.gov. All information provided in this presentation is based on information available to DURECT as of the date hereof, and DURECT assumes no obligation to update this information as a result of future events or developments, except as required by law.


Company Highlights

Larsucosterol: Phase 3-ready, potential life-saving treatment for alcohol-associated hepatitis (AH)

- **Compelling results from 307-patient placebo-controlled Phase 2b AHFIRM trial**
 - Improvement in 90-day survival and transplant-free survival
 - Well tolerated with no drug-related toxicities
- **Preparing to initiate 200-patient Phase 3 trial in U.S. with 90-day survival endpoint – data expected within 2 years of trial initiation**
 - Type B meeting with FDA resulted in agreement on key aspects of trial design
 - Single pivotal trial required for NDA submission
- **Granted Breakthrough Therapy and Fast Track designations by the FDA**
 - Breakthrough Therapy designation allows for frequent interactions with FDA and rolling NDA submission
- **Significant unmet need: >\$1 billion peak sales potential in U.S.¹**
 - 90-day mortality rate of ~30%²
 - No approved therapy for AH

¹ Market Research – ClearView Analysis 2023

² Hughes E, Hopkins LJ, Parker R. 2018. *PLOS ONE*, 13(2): e0192393



Larsucosterol
Potential in
Alcohol-associated Hepatitis

Larsucosterol: Our Lead Epigenetic Modulator Program

Phase 3-ready in AH with novel mechanism of action

Potent DNMT Modulator

- Inhibition of DNMT-1, 3a & 3b aligns with AH biology
- Supports investigating larsucosterol for the treatment of multiple acute organ injuries and chronic liver diseases

Positive Effects on Key Cellular Functions

- Stabilizes mitochondria
- Reduces lipotoxicity
- Regulates inflammation and stress response
- Promotes cell survival

Demonstrated Clinical Efficacy and Safety


- Completed Phase 2b AHFIRM trial showed compelling efficacy signal in AH patients
- Well tolerated at all doses
- >500 subjects dosed in multiple completed Phase 1 and Phase 2 studies

What is Alcohol-associated Hepatitis?

90-day mortality rate: ~30%¹

- Life-threatening form of alcohol-associated liver disease
- Can occur in individuals who chronically misuse alcohol; frequently manifests after increased consumption
- Excessive drinking can cause significant – but reversible – liver impairment leading to hepatocyte death
- Characterized by jaundice and severe multi-system inflammation – indicative of SIRS (Systemic Inflammatory Response Syndrome)²
- SIRS may progress to multi-organ failure and ultimately death

**Neutrophilic
Inflammation** **Fat Globules**



**Ballooning
Degeneration**

¹ Hughes E, Hopkins LJ, Parker R. 2018. *PLOS ONE*, 13(2): e0192393

² Crabb DW, Im GY, Szabo G, Mellinger JL, Lucey MR. 2020. *Hepatology* 71: (1) 306-333

Current Treatments for AH are Inadequate with No Approved Therapies

Corticosteroids


- Used as first-line treatment despite lack of demonstrated survival benefit
- Only 25% to 45% of patients are eligible for corticosteroids due to well known complications and contraindications^{1,2,3,4,5,6}

Liver Transplant

- Limited availability of donated organs restricts access^{3,7,8}
 - High liver transplant costs >\$875,000
 - Requires lifetime of immunosuppression

Few Programs in Clinical Development

- Most advanced competitive clinical program is in Phase 2a
- No other randomized clinical trials have reduced 90-day mortality






Larsucosterol AHFIRM Trial

Phase 2b Trial in AH to Evaluate Safety
and Efficacy of Larsucosterol

Phase 2b AHFIRM Trial Design



Trial Overview

Enrolled 307 severe AH patients with MDF² score ≥ 32 and MELD² score 21-30

Primary endpoint: Mortality or liver transplant at 90 days

Key secondary endpoint: 90-day mortality

Global trial conducted in U.S., E.U., Australia and U.K.

Placebo included active steroids at investigators' discretion¹

¹ All patients received supportive care, which for placebo patients included methylprednisolone capsules at the investigators' discretion. In order to maintain blinding, patients in the two larsucosterol arms received matching placebo capsules if the investigator prescribed steroids.

² Maddrey's Discriminant Function (MDF); Model for End-Stage Liver Disease (MELD).

Median Baseline Characteristics by Arm – Global Population

	Placebo	Larsucosterol 30 mg	Larsucosterol 90 mg
Number of patients randomized	103	102	102
Number of patients with 90-day outcome data	102 ¹	99	101
MELD ²	25.0	24.0	25.0
MDF	61.5	57.2	63.0
Age	47.0	44.0	43.0

¹ One subject in placebo group was confirmed alive at Day 90 but transplant status unknown.

² Based on central lab MELD score.


Trial Outcomes by Arm – Global Population

	Placebo ¹	Larsucosterol 30 mg	Larsucosterol 90 mg
Global:	n=102	n=99	n=101
Deaths	25 (24.5%)	15 (15.2%)	17 (16.8%)
Transplants	4 (3.9%)	6 (6.1%)	9 (8.9%)
Alive & Transplant-free	73 (71.6%)	78 (78.8%)	75 (74.3%)
All Alive	77 (75.5%)	84 (84.8%)	84 (83.2%)
U.S.:	n=77	n=73	n=77
Deaths	21 (27.3%)	8 (11.0%)	10 (13.0%)
Transplants	4 (5.2%)	5 (6.8%)	8 (10.4%)
Alive & Transplant-free	52 (67.5%)	60 (82.2%)	59 (76.6%)
All Alive	56 (72.7%)	65 (89.0%)	67 (87.0%)


¹ One subject in placebo group was confirmed alive at Day 90 but transplant status unknown. One patient received a liver transplant and subsequently died.

Clinically Meaningful Trend Toward Reduced Mortality – Global (ITT)

Mortality at 90 Days
30 mg Larsucosterol vs. Placebo¹




Mortality at 90 Days
90 mg Larsucosterol vs. Placebo¹




¹ Sites with enrollment <5 patients were pooled resulting in different mortality rates for placebo compared with 30 mg and 90 mg larucosterol doses. ITT includes 5 patients with missing 90-day outcome data. The analyses were adjusted by the method of multiple imputations to account for these subjects.

Pronounced Reduction in Mortality Observed in U.S. Patients (ITT)

Mortality at 90 Days – U.S. Patients
30 mg Larsucosterol vs. Placebo¹



Mortality at 90 Days – U.S. Patients
90 mg Larsucosterol vs. Placebo¹




¹ Sites with enrollment <5 patients were pooled resulting in different mortality rates for placebo compared with 30 mg and 90 mg larucosterol doses. ITT includes 5 patients with missing 90-day outcome data. The analyses were adjusted by the method of multiple imputations to account for these subjects.

Larsucosterol Was Well-Tolerated

Numerically fewer TEAEs in both 30 mg and 90 mg arms compared with placebo

No meaningful difference in serious AEs and none attributed to larsucosterol




Key Learnings from AHFIRM Trial

Phase 3 trial design incorporates important findings from AHFIRM trial

AHFIRM Trial Factor	Key Takeaways
1. Dose: Both doses of larsucosterol appeared to have similar efficacy and safety profiles	Conduct Phase 3 trial with 30 mg dose compared with placebo
2. Inclusion of transplants in primary endpoint: Transplants are a confounding factor	Use primary endpoint of survival
5. Regional differences: Treatment practices and disease definition vary by region	Conduct Phase 3 trial in U.S. only
4. Time to treatment: Early treatment following hospitalization is critical	Treat patients in <10 days
5. Randomization: Central randomization resulted in imbalances across arms	Randomize by site

Registrational Phase 3 Trial Design

Topline data expected within 2 years of trial initiation



A faint, light green DNA double helix structure is visible on the left side of the slide, extending from the top to the bottom. The background is a light green gradient with several diagonal, rounded rectangular bars in shades of green and teal on the right side.

Larsucosterol

Commercial Opportunity in AH

AH Imposes High Economic Burden on U.S. Healthcare System

- ~164,000 U.S. hospitalizations in 2021¹
- Incidence may result in ~300K hospitalizations by 2034² based on historical yearly growth rate of 5.5% between 2015-2019³
- Increased physician and hospital awareness of AH could result in more robust ICD-10 coding and increased recorded hospitalizations
- 88% of hospitalized AH patients are insured¹

Each hospitalization episode with AH¹:


Average hospital charges/length of stay

Died during hospitalization (2021)

~\$181,000/10 Days

Were discharged (2021)

~\$68,000/6 days



Total hospital healthcare charges per stay

¹ National Inpatient Sample 2021.

² Market Research – ClearView Analysis 2023.

³ Marlowe N et al. 2022. *Alcohol Clin Exp Res*, 46(8):1472-1481.

Larsucosterol Value Proposition Supports Blockbuster Potential

If approved, larsucosterol has the potential to become a >\$1B/year drug in the U.S. for the AH indication alone

REDUCTION IN MORTALITY

Physicians prioritize **mortality as the most important endpoint**, and nearly all found a potentially significant reduction in 90-day mortality rate clinically meaningful

HOSPITAL COST OFFSET ECONOMICS

Reducing costly **length and frequency of ICU stays** is key for offsetting drug costs and securing favorable hospital formulary inclusion


REDUCTION IN HEALTHCARE SYSTEM COST BURDEN

Hospital economics and payer stakeholders may use **reduction in 30-day readmissions** to assess impact on per-patient cost burden

Our Market Survey Indicates Physicians Are Enthusiastic About Larsucosterol


MECHANISM OF ACTION

High enthusiasm for novel, specific MOA which targets the underlying liver inflammation and degradation




CLINICAL EFFICACY

Reduction in 90-day mortality viewed as an advancement, as steroids do not show an effect on mortality past 28 days




SAFETY AND TOLERABILITY

Larsucosterol safety profile was well-received, with hundreds of patients dosed, viewed as compelling for use



DOSING AND ADMINISTRATION

Physicians saw no issues with inpatient IV doses



◆ Level of enthusiasm

Larsucosterol Intellectual Property & Regulatory Highlights

15

PATENT FAMILIES

Each include pending patent applications that, if granted, could result in protection until at least 2032 to 2044

7

ISSUED PATENTS

Seven patent families each include at least one granted patent that could provide protection until at least 2026, 2032, 2034, 2035, 2037, 2037, and 2037, respectively

**BREAKTHROUGH
THERAPY**
DESIGNATION

Designation provides intensive guidance and organizational commitment from senior FDA managers

**FAST
TRACK**
DESIGNATION

FDA designation for larsucosterol in the U.S. indicates the EMA could grant similar regulatory recognition in the EU

Larsucosterol – Positioned for Success in AH

Compelling AHFIRM Mortality Results

- Compelling outcome on key secondary endpoint of mortality reduction at 90 days
- Pronounced impact on mortality in U.S. population
- Breakthrough Therapy and Fast Track designations

Clinical Safety

- Well tolerated, no drug-related toxicities
- Numerically fewer TEAEs in AHFIRM
- No serious AEs in AHFIRM attributed to larsucosterol
- More than 500 patients dosed in multiple Phase 1 and 2 trials

>\$1 Billion Peak Sales Potential for AH in U.S.¹

- ~164,000 annual hospitalizations in U.S.²
- ~\$10 billion annual direct hospital charges in U.S.²
- No approved therapy
- Potential patent protection through at least 2044

Preparing to initiate Phase 3 trial with topline data expected within 2 years of initiation

¹ Market Research – ClearView Analysis 2023.

² National Inpatient Sample 2021.

Financial Overview

Cupertino, CA headquarters

Nasdaq	DRRX
Shares O/S	31.0 MM ¹
Cash & Cash Equivalents	\$15.1 MM ²
Debt	\$ – MM ²



¹ As of November 11, 2024

² As of September 30, 2024. Pro forma for sale of ALZET product line and repayment of Oxford term loan.

Company Highlights

Larsucosterol: Phase 3-ready, potential life-saving treatment for alcohol-associated hepatitis (AH)


- **Compelling results from 307-patient placebo-controlled Phase 2b AHFIRM trial**
 - Improvement in survival and transplant-free survival
 - Well tolerated with no drug-related toxicities
- **Preparing to initiate 200-patient Phase 3 trial in U.S. with 90-day survival endpoint – data expected within 2 years of trial initiation**
 - Type B meeting with FDA resulted in agreement on key aspects of trial design
 - Single pivotal trial required for NDA submission
- **Granted Breakthrough Therapy and Fast Track designations by the FDA**
 - Breakthrough Therapy designation allows for frequent interactions with FDA and rolling NDA submission
- **Significant unmet need: >\$1 billion peak sales in U.S.¹**
 - 90-day mortality rate of ~30%²
 - No approved therapy for AH

¹ Market Research – ClearView Analysis 2023

² Hughes E, Hopkins LJ, Parker R. 2018. *PLOS ONE*, 13(2): e0192393

Appendix


Mechanism of Action Leverages Epigenetics to Impact Disease



- 1 Epigenetic Dysregulation in AH Patients**
Aberrant DNA hypermethylation is associated with many diseases including severe AH
- 2 Epigenetic Regulators Modulate Gene Expression**
DNA methyltransferases (DNMTs) are one such regulator that add methyl groups to certain regions of DNA, generally reducing gene expression
- 3 Larsucosterol Inhibits DNMTs**
By inhibiting DNMTs (1, 3a, & 3b), larsucosterol reduces DNA hypermethylation, which modulates important cell signaling pathways


Inhibition of DNMT-1, 3a & 3b Aligns with AH

Liver samples from patients with severe AH have increased expression of DNMT-1 & 3a




Numerical Improvement in Transplant-free Survival (TFS) – Global (ITT)

Win Probability^{1,2} at 90 Days
30 mg Larsucosterol vs. Placebo



Win Probability^{1,2} at 90 Days
90 mg Larsucosterol vs. Placebo




¹ Primary endpoint was analyzed using a hierarchical assessment of patient outcomes to calculate a win probability for each of the 30 mg and 90 mg doses of larsucosterol compared with placebo. Win probability was calculated on the hierarchy of alive and transplant-free being superior to transplant and death, and transplant being superior to death. Comparisons of the same outcome were included in the denominator as ties.


² Intent-to-treat (ITT) includes 5 patients with missing 90-day outcome data. The analyses were adjusted by the method of multiple imputations to account for these subjects.

TFS – U.S. Patients (ITT)

Win Probability^{1,2} at 90 Days
30 mg Larsucosterol vs. Placebo



Win Probability^{1,2} at 90 Days
90 mg Larsucosterol vs. Placebo



¹ Primary endpoint was analyzed using a hierarchical assessment of patient outcomes to calculate a win probability for each of the 30 mg and 90 mg doses of larsucosterol compared with standard of care. Win probability was calculated on the hierarchy of alive and transplant-free being superior to transplant and death, and transplant being superior to death. Comparisons of the same outcome were included in the denominator as ties.

² Intent-to-treat (ITT) includes 5 patients with missing 90-day outcome data. The analyses were adjusted by the method of multiple imputations to account for these subjects.

AHFIRM Results by Region

	Placebo	Larsucosterol 30 mg	Larsucosterol 90 mg
France/Belgium:			
Number of subjects with 90-day outcome data	7	11	8
Deaths (%)	1 (14.3%)	4 (36.4%)	4 (50.0%)
Transplants (%)	0	1 (9.1%)	1 (12.5%)
Alive & Transplant-free (%)	6 (85.7%)	6 (54.6%)	3 (37.5%)
UK:			
Number of subjects with 90-day outcome data	1	5	2
Deaths (%)	1 (100.0%)	0	1 (50.0%)
Transplants (%)	0	0	0
Alive & Transplant-free (%)	0	5 (100.0%)	1 (50.0%)
Australia:			
Number of subjects with 90-day outcome data	17	10	14
Deaths (%)	2 (11.8%)	3 (30.0%)	2 (14.3%)
Transplants (%)	0	0	0
Alive & Transplant-free (%)	15 (88.2%)	7 (70.0%)	12 (85.7%)
U.S.¹:			
Number of subjects with 90-day outcome data	77	73	77
Deaths (%)	21 (27.3%)	8 (11.0%)	10 (13.0%)
Transplants (%)	4 (5.2%)	5 (6.8%)	8 (10.4%)
Alive & Transplant-free (%)	52 (67.5%)	60 (82.2%)	59 (76.6%)

¹ Excludes 5 subjects with missing 90-day outcome data.